



# MACHAKOS UNIVERSITY

University Examinations 2019/2020 Academic Year

SCHOOL OF PURE AND APPLIED SCIENCES

DEPARTMENT OF MATHEMATICS AND STATISTICS

.....YEAR ..... SEMESTER SPECIAL /SUPPLEMENTARY EXAMINATION FOR

BACHELOR OF SCIENCE IN ACTUARIAL SCIENCE

SAC 100: PRINCIPLES OF ACTUARIAL SCIENCE

DATE:

TIME:

---

**INSTRUCTIONS: Attempt Question ONE and any other TWO questions.**

## QUESTION ONE (30 MARKS)

- a) Define the following terms:
- A differential Equation. (1 mark)
  - Half-life. (1 mark)
  - Doubling time. (1 mark)
  - Equilibrium state. (1 mark)
  - Interactions. (1 mark)
- b) Explain two types of growth rates. (2 marks)
- c) Find both equilibrium solutions of  $\frac{ds}{dt} = 2s^2 + s - 15$  and determine their stability. (4 marks)
- d) Find the particular solution to  $\frac{dX}{dt} = \frac{3X + 5}{2}$  satisfying  $X(0) = 7$ . (4 marks)
- e) Verify that a fixed point of  $Y_{n+1} = \frac{3Y_n^2 - 6Y_n + 8}{4}$  is  $Y_{eq} = 2$  and find the other fixed point.  
Determine the stability of both fixed points. (6 marks)
- f) List three applications of Logistic Models. (3 marks)
- g) Solve  $Q_{n+2} = 5Q_{n+1} - 6Q_n$  with  $Q_0 = 0, Q_1 = 2$ . (6 marks)

## QUESTION TWO (20 MARKS)

- a) The interaction between a host species and a parasite species is modelled by the following pair of differential equations

$$P'(t) = 7P(t) - 4Q(t)$$

$$Q'(t) = 5P(t) - 2Q(t)$$

Where  $P(t)$  and  $Q(t)$  give the populations of the host and parasites at time  $t$  respectively.

The time is measured in years.

- i. Find the general solutions for  $P(t)$  and  $Q(t)$ . (4 marks)
  - ii. Find the particular solution for  $P(0) = 1500$  and  $Q(0) = 2000$  and use it to determine the long term behavior of both species. (4 marks)
  - iii. Assume that  $P(0)$  and  $Q(0)$  are both positive and let  $R$  represent the average number of parasites per host when  $t = 0$ . For which range of values of  $R$  do both species survive indefinitely. (3 marks)
- b) Determine the equilibrium and stability of  $\frac{dp}{dt} = \cos Q$ . (2 marks)

Hence:

- i. Sketch the growth rate curve (Phase diagram), (4 marks)
- ii. Sketch the general solution. (3 marks)

## QUESTION THREE (20 MARKS)

- a) Consider the prey predator system below, where  $x$  represents the prey and  $y$  the predators.

$$\frac{dx}{dt} = 3x - 9xy$$

$$\frac{dy}{dt} = -y + 4xy$$

$$\text{ii. } \frac{dx}{dt} = 7x - \frac{1}{3}xy$$

$$\frac{dy}{dt} = -y + \frac{1}{4}xy$$

- i. In which system does the prey reproduce more quickly when there are no predators (justify your answer)? (2 marks)
- ii. In which system are the predators more successful at catching prey (justify your answer)? (2 marks)

- iii. Modify the first model (*i.*) in such a way that it includes the effect of hunting of predator at a rate  $\alpha = 0.2$  proportional to the number of predator. (1 mark)
- iv. Suppose in the second model (*ii.*) that in the absence of predator, the prey population grows logistically with a carrying capacity of 80. Write the system which takes into account the above assumption. (2 marks)
- b) A public health campaign causes the contagiousness of a disease to decay exponentially. The spread of the epidemic can now be modelled by the equation  $\frac{dp}{dt} = re^{-at} p(1-p)$  where  $p$  represents the fraction of the population with the disease,  $r$  and  $a$  are both positive constants and  $t$  is the time in days.
- i. Use separation of variables and partial fractions to determine the general solution to the above differential equation. (5 marks)
- ii. If  $r = 0.2$ ,  $a = 0.04$  and  $p(0) = 0.1$  calculate the time  $t$  when half of the population is infected. (3 marks)
- iii. Using the same parameters given above, calculate the fraction of the population that is infected when  $t = 40$ . (3 marks)
- iv. Let  $R(t) = \frac{p(t)}{1-p(t)}$  be the ratio of infected to uninfected individuals. Use the general solution to show that  $\lim_{t \rightarrow \infty} R(t) = R(0)e^{r/a}$ . (2 marks)

#### QUESTION FOUR (20 MARKS)

- a) Consider the one parameter family model described by the equation  $\frac{dy}{dt} = y^3 + \alpha y + y$
- i. Locate the bifurcation value and describe the bifurcation that takes place. (4 marks)
- ii. Draw the bifurcation diagram. (3 marks)
- b) Consider a population described by the differential equation  $\frac{dy}{dt} = y^2 - 4y + 2$ .
- i. Find the equilibrium points, their stability and draw the phase diagram. (4 marks)
- ii. Describe the long-term behavior of the population with the given initial population:
- i.  $y(0) = 2$  (2 marks)

- ii.  $y(0) = 5$  (2 marks)
- c) Find the particular solution to  $\frac{dx}{dt} = 0.1x(4-x)$ , where  $x(0) = 5$ . (5 marks)

**QUESTION FIVE (20 MARKS)**

- a) The interaction of two biochemical reagents is monitored at regular intervals and can be modelled by the following pair of coupled recurrence relations

$$X_{n+1} = \frac{6}{5}X_n + \frac{7}{10}Y_n$$

$$Y_{n+1} = \frac{3}{10}X_n + \frac{4}{5}Y_n$$

Where  $X_n$  and  $Y_n$  denote the amount of each reagent after  $n$  observations.

- i. Show that the characteristic equation is  $r^2 - 2r + \frac{3}{4} = 0$ . (2 marks)
- ii. Find the general solution for  $X_n$  and  $Y_n$ . (4 marks)
- iii. Find the particular solution when  $X_0 = 900$  and  $Y_0 = 100$ . (3 marks)
- iv. Show that as  $n \rightarrow \infty$  the ratio  $X_n/Y_n$  tends to  $7/3$ . (1 mark)
- v. Show that the limit in the answer to part (d) does not depend on the initial amounts. (2 marks)
- b) Given  $X_{n+1} = \frac{1}{6}X_n^2(5 - X_n)$ . Find:
- i. Fixed points. (2 marks)
- ii. Equilibrium condition and stability. (3 marks)
- iii. Approximate general solution. (3 marks)