



# MACHAKOS UNIVERSITY

University Examinations 2021/2022

SCHOOL OF PURE AND APPLIED SCIENCES

DEPARTMENT OF PHYSICAL SCIENCES

SECOND YEAR SUPPLEMENTARY/SPECIAL EXAMINATION FOR

BACHELOR OF EDUCATION SCIENCE (SPECIAL NEEDS)

BACHELOR OF SCIENCE MATHEMATICS

BACHELOR OF EDUCATION (SCIENCE)

SCH 201: CHEMICAL THERMODYNAMICS

DATE: 16/03/2022

TIME: 8:30-10:30 AM

---

## INSTRUCTIONS:

- The paper consists of **two** sections.
- Section **A** is **compulsory** (30 marks).
- Answer any **two** questions from section **B** (each 20 marks).

## SECTION A

### QUESTION ONE (30 MARKS)

- a) State and give a mathematical expression for the first law of thermodynamics (2 marks)
- b) Classify the following properties of matter as either extensive or intensive (3 marks)
- Pressure
  - Density
  - Enthalpy
- c) Explain why  $C_p > C_v$  (3 marks)
- d) Classify the following as isolated, closed or open system (4 marks)
- Nitrogen and hydrogen reacting in a sealed tube to form ammonia
  - A glass vial containing Sulphuric acid is broken inside water in a beaker
  - Vacuum flask
  - The Earth

- e) Two Carnot engines operate with a temperature difference of 300 K between their hot and cold reservoirs. If engine A's hot reservoir is maintained at 1200 K, while that of engine B is maintained at 750 K, determine the most efficient engine. (3 marks)
- f) Using second law of thermodynamics show that for a cyclic engine with no cold reservoir the conversion of all heat to equivalent work is non-spontaneous. (3 marks)
- g) Calculate the change in free energy when 11.21dm<sup>3</sup> of a perfect gas at 0°C and 750mmHg pressure expands isothermally until its pressure is 180mmHg. (3 marks)
- h) The Critical temperature of methane and carbon dioxide are -81.9°C and 31.1°C respectively. Determine which of these gases has stronger intermolecular forces and why? (3 marks)
- i) Using  $G = H - TS$ , and  $dH - vdp - tds \leq 0$ , derive an alternative criterion for spontaneity at constant temperature and pressure (3 marks)
- j) Determine the entropy of the following reaction using the information below. (3 marks)

| Substance                        | S°(Jmol <sup>-1</sup> K <sup>-1</sup> ) |
|----------------------------------|---|
| C <sub>2</sub> H <sub>2(g)</sub> | 200.82                                  |
| H <sub>2(g)</sub>                | 130.59                                  |
| C <sub>2</sub> H <sub>4(g)</sub> | 219.45                                  |

$$\text{C}_2\text{H}_2(\text{g}) + \text{H}_2(\text{g}) \longrightarrow \text{C}_2\text{H}_4(\text{g})$$

## SECTION B

### QUESTION TWO (20 MARKS)

- a) State the Hess's law.
- b) Calculate the enthalpy of the reaction,  $\text{C}_2\text{H}_4(\text{g}) + \text{H}_2(\text{g}) \longrightarrow \text{C}_2\text{H}_6(\text{g})$  at 298 k using the information given below. (6 marks)
- $$\text{C}_2\text{H}_4(\text{g}) + 3\text{O}_2(\text{g}) \longrightarrow 2\text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g}) \quad \Delta H = -1395 \text{ kJ}$$
- $$\text{C}_2\text{H}_6(\text{g}) + \frac{7}{2}\text{O}_2(\text{g}) \longrightarrow 2\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{g}) \quad \Delta H = -1550 \text{ kJ}$$
- $$\text{H}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \longrightarrow \text{H}_2\text{O}(\text{g}) \quad \Delta H = -243 \text{ kJ}$$
- c) The volume of a sample of an ideal monoatomic gas at 0°C is 44.83 litres. To what volume must the gas be compressed adiabatically to attain a temperature of 30°C? (6 marks)
- d) State and explain the physical concept of entropy and its variation with temperature. (6 marks)

### QUESTION THREE (20 MARKS)

- a) Sample of 1 mole of silver at 150°C is placed in contact with 1mole of silver at 0°C forming an isolated system at constant pressure. Assuming the specific heat capacity of silver is 28.85J/mol k, Calculate
- i) The final temperature of both the silver samples. (4 marks)
  - ii)  $\Delta S$  for the hot silver. (2 marks)
  - iii)  $\Delta S$  for the cold silver. (2 marks)
  - iv) The total  $\Delta S$  of the system. (1 mark)
  - v) Is the process above spontaneous. (1 mark)
- b) Provide an expression for work done by a gas as it expands by volume  $\Delta V$  against an external pressure  $P_{\text{ext}}$ . (2 marks)
- c) Give an expression for the change in internal energy of the system as it expands by volume  $\Delta V$  against an external pressure  $P_{\text{ext}}$  on absorbing  $q$  joules of heat. (3 marks)
- d) A sample of gas changes in volume from 6 Litres to 8 Litres against an external pressure of 2.5 atm and simultaneously absorbs 1000 J of heat. What is the change in the internal energy of the system? (5 marks)

### QUESTION FOUR (20 MARKS)

- a) A 4 L sample of hydrogen gas at 15 atm and 25°C is allowed to expand to a final pressure of 1atm. Calculate  $q$ ,  $w$ ,  $\Delta U$  and  $\Delta H$  if the gas expands isothermally and reversibly. (10 marks)
- b) A certain gas expands from 400 mL to 1200 mL very slowly at 25°C. If there is one mole of the gas in the chamber, calculate the work involved when the gas behaves ideally. (5 marks)
- c) Given the following standard enthalpies of formation, calculate the standard enthalpy of combustion ( $\Delta H^\circ_{\text{combustion}}$ ) of benzene at 298 k. (5 marks)

| Compound                      | $\Delta H^\circ_f$ (kJ/mol k) |
|-------------------------------|-------------------------------|
| CO <sub>2</sub>               | -393.51                       |
| C <sub>6</sub> H <sub>6</sub> | +48.7                         |
| H <sub>2</sub> O              | -285.83                       |

### QUESTION FIVE (20 MARKS)

- a) The standard molar enthalpy of formation of ammonia is  $-56.25 \text{ kJ/mol}$ . using the heat capacity data given below, calculate the standard molar heat of formation of ammonia at  $1000 \text{ K}$ . (10 marks)

$$C_p \text{N}_2(\text{g}) \text{ J/mol. } k=26.98+(5.912 \times 10^{-3}) T$$

$$C_p \text{NH}_3(\text{g}) \text{ J/mol. } k=25.89+(32.58 \times 10^{-3}) T$$

$$C_p \text{H}_2(\text{g}) \text{ J/mol. } k=29.07-(0.837 \times 10^{-3}) T$$

- b) Describe the boundary conditions in;
- i) Open system (2 marks)
  - ii) Closed system (2 marks)
  - iii) Isolated system (2 marks)
- c) Define the following terms
- i) Specific heat capacity (2 marks)
  - ii) Molar heat capacity (2 marks)

*END*