



MACHAKOS UNIVERSITY

University Examinations 2018/2019

SCHOOL OF PURE AND APPLIED SCIENCES

DEPARTMENT OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE

THIRD YEAR SECOND SEMESTER EXAMINATION FOR

BACHELOR OF SCIENCE (MATHEMATICS AND COMPUTER SCIENCE)

BACHELOR OF SCIENCE (STATISTICS AND PROGRAMMING)

BACHELOR OF SCIENCE (MATHEMATICS)

BACHELOR OF EDUCATION (SCIENCE)

BACHELOR OF EDUCATION (ARTS)

SMA 365 : DESIGN AND ANALYSIS OF SAMPLE SURVEYS

DATE: 15/4/2019

TIME: 2:00 – 4:00 PM

Instructions to the Candidate:

1. Answer **Question 1** and any other **two** questions.
 2. Out of the **three** questions answered, each question must start on a new page.
 3. You need a Scientific Calculator and Statistical Tables for this paper
-
1. a) Explain the importance of a *pre-test* in the collection of statistical data in a survey. (6 marks)
 - b) With the aid of an example in each case, differentiate between each of the following terms as used in the collection of statistical data.
 - (i) Observational unit and sampling unit;
 - (ii) Census and sample survey. (6 marks)
 - c) Given an SRS-WOR with a sample mean $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ and a population mean $\bar{X} = \frac{1}{N} \sum_{i=1}^N x_i$.
Show that the sample mean $\bar{x}$ is an unbiased estimator of the population mean $\bar{X}$. (4 marks)

- d) Given a stratified random sample with a stratified mean given by $\bar{X}_{st} = \frac{1}{N} \sum_{i=1}^m N_i \bar{x}_i$ and a population mean given by $\bar{X} = \frac{1}{N} \sum_{i=1}^m X_i$, show that the stratified mean $\bar{X}_{st}$ is an unbiased estimator of the population mean $\bar{X}$. (4 marks)

- e) Using a population of size $N = 5$ comprising the values 24, 21, 18, 27, 15, from which a random sample of size $n = 3$ is to be selected, determine all the possible samples of size 3 which can be drawn from the population.

Hence using the selected samples, prove that the sample mean $\bar{x}$ is an unbiased estimator of the population mean $\bar{X}$. (10 marks)

2. a) Explain the term *multi-stage sampling* as used in the collection of statistical data. (2 marks)

- b) Explain the circumstance under which each of the following sampling techniques is suitable in the collection of statistical data.

- (i) Simple random sampling;
- (ii) Stratified random sampling. (5 marks)

- c) A survey was conducted to determine the number of mobile phone SIM cards used by households among Kenyan urban residents. A simple random sample of 25 households was drawn from a residential estate comprising 800 households. The number of phone SIM cards owned in a household in the sample was as follows:

8, 6, 5, 7, 9, 4, 8, 5, 4, 6, 9, 3, 8, 9, 6, 5, 7, 9, 6, 4, 8, 4, 6, 5, 9.

- i) Estimate the total number of phone SIM cards owned by the households in the residential estate; (3 marks)
- ii) Determine the 95% confidence interval for the population mean of the number of phone SIM cards owned by residents in the residential estate; (5 marks)
- iii) Determine the 95% confidence interval for the population total of the number of phone SIM cards owned by residents in the estate. (5 marks)

3. a) Explain the term *questionnaire* as used in the collection of statistical data. (2 marks)

- b) Explain the **three** errors which may occur through sampling in a sample survey, illustrating each with an example from a real life situation. (9 marks)

- c) Given an SRS-WOR with a sample variance given by $s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$ and its corresponding population variance given by $S^2 = \frac{1}{N-1} \sum_{i=1}^N (x_i - \bar{X})^2$, show that the sample variance s^2 is an unbiased estimator of the population variance S^2 . (9 marks)

4. a) Explain the term *sampling frame* as used in a sample survey, giving *two* examples from real life situations. (4 marks)
- b) Given a two-stage sample in which the sample mean is given by $\bar{\bar{x}} = \frac{1}{n m} \sum_{i=1}^n \sum_{j=1}^m x_{ij}$ and its corresponding population mean is given by $\bar{\bar{X}} = \frac{1}{N M} \sum_{i=1}^N \sum_{j=1}^M x_{ij}$, show that the sample mean $\bar{\bar{x}}$ is an unbiased estimator of the population mean $\bar{\bar{X}}$. (6 marks)
- c) A survey is to be conducted among patients in a county hospital, with the patients categorised as children, teenagers, middle age and old age. A sample of size $n = 20$ is to be selected from a population of size $N = 800$. Select an *actual sample* procedurally stating clearly the range of your sampling frame and the selected sampling units using each of the following sampling techniques:
- Simple random sampling; (6 marks)
 - Systematic sampling; (4 marks)
5. A survey was conducted to determine the number of cattle kept in homesteads among a pastoralist community in Kenyan. A stratified random sample of size 140 homesteads was drawn from a target population of size 2,800 homesteads. This target population is divided into four strata as shown below:

Stratum	A	B	C	D
Stratum size	800	900	600	500
Stratum sample mean	64	75	80	74
Stratum standard deviation	4	5	8	3

- a) Determine the sample size for each stratum if the sample is to be drawn using stratified random sampling technique with:
- proportional allocation;
 - optimum allocation with fixed cost. (10 marks)
- b) Using the sample sizes obtained under proportional allocation, determine the 95% confidence limits for each of the following measures about the pastoralists:
- the population mean of the number of cattle kept in a homestead; (5 marks)
 - the population total of the number of cattle kept by the pastoralists. (5 marks)