



MACHAKOS UNIVERSITY

University Examinations 2019/2020 Academic Year

SCHOOL OF PURE AND APPLIED SCIENCES

DEPARTMENT OF MATHEMATICS AND STATISTICS

FOURTH YEAR SPECIAL /SUPPLEMENTARY EXAMINATION FOR

BACHELOR OF SCIENCE (STATISTICS AND PROGRAMMING)

BACHELOR OF SCIENCE (MATHEMATICS AND COMPUTER SCIENCE)

BACHELOR OF SCIENCE (MATHEMATICS)

BACHELOR OF EDUCATION (SCIENCE)

SMA 463 TIME SERIES ANALYSIS

DATE: 20/1/2021

TIME: 2.00-4.00 PM

INSTRUCTIONS TO CANDIDATES

Answer Question ONE and ANY OTHER TWO

QUESTION ONE (30 MARKS)

- a) Explain briefly the four components of a time series. (8 marks)
- b) Giving an example, explain one objective of time series analysis. (2 marks)
- c) Differentiate between strict stationarity and weak stationarity. (4 marks)
- d) Let $\{X_t\}, t = 0, \pm 1, \pm 2, \dots$ be a stochastic process given by $X_t = a + bt + e_t$ where e_t is a sequence of independent random variables distributed with mean $\mu = 0$ and variance σ^2 as $e_t \sim N(0, \sigma^2)$. Show that $\nabla X_t = X_t - X_{t-1}$ is second order (weakly) stationary. (8 marks)
- e) Explain briefly the importance of the periodogram in spectral analysis. (2 marks)
- f) Consider the first-order moving average process defined by the equation
$$X_t = Z_t + \theta Z_{t-1}, t = 0, \pm 1, \dots$$
 where $\{Z_t\} \sim WN(0, \sigma^2)$, θ is a real valued constant and $EX_t = 0$.
Determine;
 - i. $\gamma_x(t + h, t)$ (2 marks)
 - ii. The spectral density of the process. (4 marks)

QUESTION TWO (20 MARKS)

- a) Explain briefly the procedure you would adopt in the analysis of a time series. (4 marks)
- b) Describe one method each of
- Eliminating the effect of trend from a time series.
 - Measuring seasonal variations.
 - In measuring seasonal variation, can cyclical and erratic influences be eliminated? How? (5 marks)
- c) The table below gives the world production of gold (in crore ounces). Fit a straight line trend by the method of least squares. (6 marks)

Year	Production
1991	12.7
1992	10.1
1993	13.0
1994	13.2
1995	12.6
1996	14.2
1997	13.7

- d) Suppose $Y_t = \beta_0 + \beta_1 t + X_t$, where $\{X_t\}$ is a zero-mean stationary series with autocovariance function γ_k and β_0 and β_1 are constants. Show that $\{Y_t\}$ is not stationary but that $\nabla Y_t = Y_t - Y_{t-1}$ is stationary. (5 marks)

QUESTION THREE (20 MARKS)

- a) Define an ARMA (p, q) process. (2 marks)
- b) Explain briefly the three properties of the autocorrelation function. (3 marks)
- c) Consider two MA(1) processes P and Q given by

$$P: X_t = Z_t + \theta Z_{t-1}$$


$$Q: X_t = Z_t + \frac{1}{\theta} Z_{t-1}$$

Obtain the autocorrelation function of each of the processes. Is a moving average process uniquely identified from a given autocorrelation function? Explain. (7 marks)

- d) Given that Z_t is iid noise with zero mean and variance σ^2 , find the autocorrelation function for the stationary process defined by $Y_t = 5 + Z_t - \frac{1}{2}Z_{t-1} + \frac{1}{4}Z_{t-2}$ (8 marks)

QUESTION FOUR (20 MARKS)

- a) Describe the important characteristics of the autocorrelation function for the following models.
- MA(1)
 - AR(1)
 - ARMA(1,1) (3 marks)
- b) Given the following ACF plots, explain with reason the generating process. (2 marks)



- c) Consider the second order autoregressive process given by $Y_t = \alpha_1 Y_{t-1} + \alpha_2 Y_{t-2} + e_t$ where e_t is a purely random process with mean zero and variance σ^2 .
- Derive the Yule-Walker equation for the process and its general solution. Find the values of α_1, α_2 , such that the process is stationary. (8 marks)
 - If $\alpha_1 = 1/3$ and $\alpha_2 = 2/9$, show that the autocorrelation function of X_t is given by

$$\rho(h) = \frac{16}{21} \left(\frac{2}{3}\right)^{|h|} + \frac{5}{21} \left(-\frac{1}{3}\right)^{|h|}, \quad h = 0, \pm 1, \pm 2, \dots$$
 (7 marks)

QUESTION FIVE (20 MARKS)

- a) Define the spectral density function of a time series X_t and show that it can be expressed in terms of cosines. (4 marks)
- b) Find the spectral density function of an AR (1) process given by $X_t = \alpha X_{t-1} + e_t$ (6 marks)
- c) Determine if the following ARMA process is causal and if it is invertible. (5 marks)

$$X_t + 0.2 X_{t-1} - 0.48 X_{t-2} = Z_t$$

- d) Consider the random walk $S_t, t = 0, 1, 2, \dots$ obtained by setting $S_0 = 0$ and $S_t = X_1 + X_2 + \dots + X_t$, for $t = 1, 2, \dots$ where $\{X_t\}$ is iid noise with $E(X_t) = 0$ and $E(X_t^2) < \infty$.

Determine;

- $E(S_t)$ (1 mark)
- $E(S_t^2)$ (2 marks)
- $\gamma_s(t+h, t)$ (2 marks)