



MACHAKOS UNIVERSITY
University Examinations 2020/2021

SCHOOL OF PURE AND APPLIED SCIENCES

DEPARTMENT OF PHYSICAL SCIENCES

THIRD YEAR SUPPLEMENTARY/SPECIAL EXAMINATION FOR
BACHELOR OF SCIENCE (MATHS),
BACHELOR SCIENCE (TELECOMMUNICATION & IT),
BACHELOR SCIENCE (APPLIED PHYSICS AND TECHNOLOGY,
BACHELOR EDUCATION (SCIENCE)
SPH 203: THERMAL PHYSICS 1

DATE:

TIME:

INSTRUCTIONS:

- The paper consists of **two** sections.
- Section **A** is **compulsory** (30 marks).
- Answer any **two** questions from section **B** (each 20 marks).

The following constants may be useful

The following constants may be useful, n is number of moles, Universal gas constant $R = 8.31 \text{ J/mol.K}$, specific heat capacity of gold and water is $129 \text{ J/kg.}^\circ\text{C}$ and $4186 \text{ J/kg.}^\circ\text{C}$ respectively, Latent heat of fusion of ice $3.33 \times 10^5 \text{ J/kg}$, specific heat capacity of argon $c_p = 20.5$ and $c_v = 12.5$, $1 \text{ atm} = 1.01 \times 10^5 \text{ N/m}^2$. Coefficient of linear expansion of copper is $17 \times 10^{-6} (^\circ\text{C})^{-1}$, for steel is $11 \times 10^{-6} (^\circ\text{C})^{-1}$, for aluminum $24 \times 10^{-6} (^\circ\text{C})^{-1}$

SECTION ONE

QUESTION ONE.

- a) Some water trapped a volcano undergoes heating and blows off the volcano in a big explosion, give an explanation why this happens. (1 marks)
- b) The lowest temperature -273.15 cannot be measured directly. Explain how it can be inferred with aid of well labeled diagram. (4 marks)
- c) A man is seated on a $10\text{ }^{\circ}\text{C}$ concrete floor in a police cell, insulation provided by his clothing is 0.5 cm thick with a coefficient of thermal conductivity of $0.09\text{ J/s.m.}^{\circ}\text{C}$ and the area of contact is 0.10 m^2 . If his body temperature is $37\text{ }^{\circ}\text{C}$, compute his rate of heat loss. (4 marks)
- d) A person with a skin temperature of $33\text{ }^{\circ}\text{C}$ is in a room at $24\text{ }^{\circ}\text{C}$. What is the net rate of heat transfer by radiation? (4 marks)
- e) 3 moles of Hydrogen gas initially at $10\text{ }^{\circ}\text{C}$ and at 1.00 atm expands to twice its initial volume isothermally in one case and adiabatically in another. Calculate the work done during the
 - i) Isothermal process (5 marks)
 - ii) Adiabatic process (5 marks)
- f) A fluorescent tube of cross sectional area 10 cm^2 and length 100 m contains argon gas at $20\text{ }^{\circ}\text{C}$. It is being heated at a rate of 50 J/s from 7.00 pm to 10 pm at constant atmospheric pressure. (assume the process is adiabatic)
 - i) Find total heat energy absorbed by the gas (2 marks)
 - ii) Determine the number of moles of argon gas in the tube (2 marks)
 - iii) Find the temperature rise? (3 marks)

QUESTION TWO.

A cylinder with a movable piston contains 0.016 mole of helium. A researcher expands the gas via the process illustrated in figure 1.

- a) To achieve this, does she need to heat the gas or cool it? (2 marks)
- b) From the graph read the initial and final volume of the gas in SI units (4 marks)
- c) Using ideal gas law find initial and final temperature of the gas in kelvin (K) (4 marks)
- d) Compute change in internal energy (ΔU) given that $\Delta U = \frac{3}{2}nR\Delta T$ (3 marks)
- e) From the graph estimate work done by the gas? (4 marks)
- f) How much heat energy must be added or removed? (3 marks)




Figure 1

QUESTION THREE.

- Explain the changes that water undergoes while changing the phase from solid to liquid at 0°C (2 marks)
- Show that thermal coefficient of volume (β) is thrice as large as the coefficient of linear expansion (α) (6 marks)
- Temperature of water in Lake Naivasha changes from 12°C to 16°C given that the thermal coefficient of volume for water is $\beta = 2.07 \times 10^{-4} (\text{C}^\circ)^{-1}$ and coefficient of linear expansion (α)
 - Estimate the fractional change in volume of water in lake Naivasha, (6 marks)
 - If initial depth of the Lake is $4.0 \times 10^3 \text{ m}$, find the change in depth. (6 marks)

QUESTION FOUR

- A plate with a circular hole (Figure 2) on the center is heated. What happens to the hole, does it become larger or smaller. (4 marks)




Figure 2

- A squared metallic plate of side L is heated from initial temperature T_i to a final temperature T_f . If the new dimension of its side is L_f , derive an equation for its fractional change in area. (8 marks)

- c) A steel ring with a hole having area of 3.99 cm^2 is to be placed on an aluminum rod with cross sectional area of 4 cm^2 . Both ring and rod are initially at a temperature of 35°C . At what common temperature can the steel ring be slipped onto one end of the aluminum rod? (8 marks)

QUESTION FIVE.

- a) What mass of water at 25°C must be allowed to come to thermal equilibrium with a 3 kg gold bar at 100°C in order to lower the temperature of the bar to 50°C ? (10 marks)
- b) A 100 g cube of ice at 0°C is dropped into 1.0 kg of water that was originally at 80°C . What is the final temperature of the water after the ice has melted? (10 marks)